

# Factorization of classical characters twisted by roots of unity

Arvind Ayyer and Nishu Kumari\* (Indian Institute of Science)


## Abstract

- Fix  $t \geq 2$ ,  $n \in \mathbb{Z}^+$ . Consider the irreducible characters of representations of  $GL_{tn}$ ,  $SO_{2tn+1}$ ,  $Sp_{2tn}$  and  $O_{2tn}$  over  $\mathbb{C}$ , evaluated at elements  $\omega^k x_i$  for  $0 \leq k \leq t-1$  and  $1 \leq i \leq n$ , where  $\omega$  is a primitive  $t^{\text{th}}$  root of unity.
- Motivated by the case of  $GL_{tn}$ , considered by D. J. Littlewood (AMS press, 1950) and independently by D. Prasad (Israel J. Math., 2016).
- We characterize partitions for which the specialized irreducible character is nonzero in terms of what we call  $z$ -asymmetric partitions, where  $z$  is an integer which depends on the group.
- The non-zero character factorizes into characters of smaller classical groups.
- We also give product formulas for general  $z$ -asymmetric partitions and  $t$ -cores.
- Finally, we show that there are infinitely many  $z$ -asymmetric  $t$ -cores for  $t \geq z+2$ .


## Notations and Definitions

$X = (x_1, \dots, x_n)$  - a tuple of commuting indeterminates.  $X^j = (x_1^j, \dots, x_n^j)$ ,  $j \in \mathbb{Z}$ .  $\bar{X} = (\frac{1}{x_1}, \dots, \frac{1}{x_n})$ .

**Partition and its beta-set:**



$$\lambda = (4, 4, 2, 1) \vdash 11, \ell(\lambda) = 4, \text{rk}(\lambda) = 2$$




$$\beta(\lambda, 4) = (7, 6, 3, 1)$$


For  $\lambda = (\lambda_1, \dots, \lambda_k)$  and  $\mu = (\mu_1, \dots, \mu_j)$  partitions,  $k + j \leq 2n$ ,

$$\mu_1 + (\lambda, 0, \dots, 0, -\text{rev}(\mu)) = (\mu_1 + \lambda_1, \dots, \mu_1 + \lambda_k, \underbrace{\mu_1, \dots, \mu_j}_{2n-j-k}, \mu_1 - \mu_j, \dots, \mu_1 - \mu_2, 0).$$

The parts of the beta set congruent to  $i \pmod t$  for  $i \in [0, t-1]$ :




$$i = 0, t = 3$$




$$i = 1, t = 3$$

**$t$ -core of  $\lambda$ :** Consider  $tj + i$ ,  $0 \leq j \leq n_i(\lambda, m) - 1$ ,  $0 \leq i \leq t-1$




$$\text{core}_3(\lambda) = (4-3, 3-2, 1-1, 0) = (1, 1).$$

**$t$ -quotient of  $\lambda$ :** For each  $i \in [0, t-1]$ , consider  $\left[ \frac{\beta^{(i)}(\lambda, m)}{t} \right]$



$$\text{quo}_3(\lambda) = (\lambda^{(0)}, \lambda^{(1)}, \lambda^{(2)}), \quad \lambda^{(0)} = (2-1, 1-0) = (1, 1), \quad \lambda^{(1)} = (2-1, 0-0) = (1), \quad \lambda^{(2)} = \emptyset.$$

**$z$ -asymmetric partition:**  $(\alpha | \alpha + z)$ ,  $z \in \mathbb{Z}$ .



$$\lambda = (4, 2, 2, 1, 1) = (3, 0 | 4, 1)$$

symplectic (1-asymmetric) 3-core

## Weyl Character Formulas

Let  $\lambda$  be a partition of length at most  $n$ .

The Schur polynomial or general linear (type A) character of  $GL_n$  indexed by  $\lambda$ :

$$s_\lambda(X) = \frac{\det_{1 \leq i, j \leq n} (x_i^{\beta_j(\lambda, n)})}{\det_{1 \leq i, j \leq n} (x_i^{n-j})}$$

The odd orthogonal (type B) character of the group  $SO(2n+1)$  indexed by  $\lambda$ :

$$\text{so}_\lambda(X) = \frac{\det_{1 \leq i, j \leq n} (x_i^{\beta_j(\lambda, n)+1/2} - \bar{x}_i^{\beta_j(\lambda, n)+1/2})}{\det_{1 \leq i, j \leq n} (x_i^{n-j+1/2} - \bar{x}_i^{n-j+1/2})}$$

The symplectic (type C) character of the group  $Sp(2n)$  indexed by  $\lambda$ :

$$\text{sp}_\lambda(X) = \frac{\det_{1 \leq i, j \leq n} (x_i^{\beta_j(\lambda, n)+1} - \bar{x}_i^{\beta_j(\lambda, n)+1})}{\det_{1 \leq i, j \leq n} (x_i^{n-j+1} - \bar{x}_i^{n-j+1})}$$

The even orthogonal (type D) character of the group  $O(2n)$  indexed by  $\lambda$ :

$$o_\lambda^{\text{even}}(X) = \frac{2 \det_{1 \leq i, j \leq n} (x_i^{\beta_j(\lambda, n)} + \bar{x}_i^{\beta_j(\lambda, n)})}{(1 + \delta_{\lambda, 0}) \det_{1 \leq i, j \leq n} (x_i^{n-j} + \bar{x}_i^{n-j})}$$

where  $\delta$  is the Kronecker delta.

For  $\ell(\lambda) \leq tn$ , let  $\sigma_\lambda \in S_{tn}$  be the permutation that rearranges the parts of  $\beta(\lambda, tn)$  such that

$$\beta_{\sigma_\lambda(j)}(\lambda, tn) \equiv q \pmod t, \quad \sum_{i=0}^{q-1} n_i(\lambda, tn) + 1 \leq j \leq \sum_{i=0}^q n_i(\lambda, tn),$$

arranged in decreasing order for each  $q \in \{0, 1, \dots, t-1\}$ .

## Schur Factorization

Theorem (D. J. Littlewood (AMS press, 1950), D. Prasad (Israel J. Math., 2016))

Let  $\lambda$  be a partition of length at most  $tn$  indexing an irreducible representation of  $GL_{tn}$  and  $\text{quo}_t(\lambda) = (\lambda^{(0)}, \dots, \lambda^{(t-1)})$ . Then the Schur polynomial  $s_\lambda(X, \omega X, \dots, \omega^{t-1} X)$  is given as follows.

1. If  $\text{core}_t(\lambda)$  is non-empty, then

$$s_\lambda(X, \omega X, \dots, \omega^{t-1} X) = 0.$$

2. If  $\text{core}_t(\lambda)$  is empty, then

$$s_\lambda(X, \omega X, \dots, \omega^{t-1} X) = \text{sgn}(\sigma_\lambda) (-1)^{\frac{n(n+1)t(t-1)}{2}} \prod_{i=0}^{t-1} s_{\lambda^{(i)}}(X^t).$$

## Factorization of other Classical Characters

Theorem (Ayyer-Kumari, [1], 2021)

Let  $\lambda$  be a partition of length at most  $tn$  indexing an irreducible representation of  $Sp_{2tn}$  and  $\text{quo}_t(\lambda) = (\lambda^{(0)}, \dots, \lambda^{(t-1)})$ . Then the  $Sp_{2tn}$ -character  $\text{sp}_\lambda(X, \omega X, \dots, \omega^{t-1} X)$  is given as follows.

1. If  $\text{core}_t(\lambda)$  is not a symplectic  $t$ -core, then

$$\text{sp}_\lambda(X, \omega X, \dots, \omega^{t-1} X) = 0.$$

2. If  $\text{core}_t(\lambda)$  is a symplectic  $t$ -core with rank  $r$ , then

$$\text{sp}_\lambda(X, \omega X, \dots, \omega^{t-1} X) = (-1)^e \text{sgn}(\sigma_\lambda) \text{sp}_{\lambda^{(t-1)}}(X^t) \prod_{i=0}^{\lfloor \frac{t-3}{2} \rfloor} s_{\mu_i}(X^t, \bar{X}^t) \times \begin{cases} \text{so}_{\lambda^{(t/2-1)}}(X^t) & t \text{ even,} \\ 1 & t \text{ odd,} \end{cases}$$

where

$$e = - \sum_{i=\lfloor \frac{t}{2} \rfloor}^{t-2} \binom{n_i(\lambda) + 1}{2} + \begin{cases} \frac{n(n+1)}{2} + nr & t \text{ even,} \\ 0 & t \text{ odd,} \end{cases}$$

and  $\mu_i = \lambda_1^{(t-2-i)} + (\lambda^{(i)}, 0, \dots, 0, -\text{rev}(\lambda^{(t-2-i)}))$ ,  $0 \leq i \leq \lfloor \frac{t-3}{2} \rfloor$ .

**Example:**  $t = 2$ ,  $n = 1$  and  $a \geq b \geq 0$ .  $\text{sp}_{(a,b)}(x, -x)$  is nonzero if and only if  $a$  and  $b$  have the same parity.

$$\text{sp}_{(a,b)}(x, -x) = \begin{cases} -\text{sp}_{(\frac{a-1}{2})}(x^2) \text{so}_{(\frac{b+1}{2})}(x^2) & a \text{ and } b \text{ are odd,} \\ \text{sp}_{(\frac{a}{2})}(x^2) \text{so}_{(\frac{b}{2})}(x^2) & a \text{ and } b \text{ are even.} \end{cases}$$

We give similar factorization results for the irreducible characters of classical groups of type B and D, namely  $O_{2tn}$  [1, Theorem 2.15] and  $SO_{2tn+1}$  [1, Theorem 2.17], where we specialize the elements as before.

## Generating Functions

The set of  $z$ -asymmetric partitions and  $z$ -asymmetric  $t$ -cores -  $\mathcal{P}_z$  and  $\mathcal{P}_{z,t}$  respectively.

Theorem (Ayyer-Kumari, [1], 2021)

For  $z \in \mathbb{Z}$ ,

$$\sum_{\lambda \in \mathcal{P}_z} q^{|\lambda|} = \prod_{k \geq 0} (1 + q^{z+1+2k}) = (-q^{z+1}; q^2)_\infty, \quad (a; q)_\infty = \prod_{j=0}^{\infty} (1 - aq^j).$$

Theorem (Ayyer-Kumari, [1], 2021)

For  $|z| \geq t-1$ , the empty partition is the only  $t$ -core in  $\mathcal{P}_{z,t}$ .

Theorem (Ayyer-Kumari, [1], 2021)

Let  $0 \leq z \leq t-2$ . Represent elements of  $\mathbb{Z}^{\lfloor \frac{t-z}{2} \rfloor}$  by  $(z_0, \dots, z_{\lfloor \frac{t-z}{2} \rfloor})$  and define  $b \in \mathbb{Z}^{\lfloor \frac{t-z}{2} \rfloor}$  by  $\vec{b}_i = t - z - 1 - 2i$ . Then there exists a bijection  $\phi: \mathcal{P}_{z,t} \rightarrow \mathbb{Z}^{\lfloor \frac{t-z}{2} \rfloor}$  satisfying  $|\lambda| = t \|\phi(\lambda)\|^2 - \vec{b} \cdot \phi(\lambda)$ , where  $\cdot$  represents the standard inner product.

**Ramanujan theta function:**

$$f(a, b) = \sum_{n=-\infty}^{\infty} a^{\frac{n(n+1)}{2}} b^{\frac{n(n-1)}{2}}.$$

Corollary (Ayyer-Kumari, [1], 2021)

Let  $p_{z,t}(m)$  be the cardinality of partitions in  $\mathcal{P}_{z,t}$  of size  $m$ . For  $0 \leq z \leq t-2$ , we have

$$\sum_{m \geq 0} p_{z,t}(m) q^m = \prod_{i=0}^{\lfloor (t-z-2)/2 \rfloor} f(q^{2i+z+1}, q^{2t-2i-z-1}).$$

## Reference

[1] A. Ayyer, N. Kumari, *Factorization of Classical characters twisted by roots of unity*, to appear in Journal of Algebra. arXiv identifier: 2109.11310.