

Dimension-free L_p estimates for maximal Riesz transforms

Błażej Wróbel,
Institute of Mathematics, Polish Academy of Sciences &
Institute of Mathematics, University of Wrocław

Based on joint work with Maciej Kucharski and Jacek Zienkiewicz

Notation:

- d - dimension of the space \mathbb{R}^d , $L^p = L^p(\mathbb{R}^d)$, $\|\cdot\|_p = \|\cdot\|_{L^p}$
- k - fixed non-negative integer
- \mathcal{H}_k - space of spherical harmonics of degree k on \mathbb{R}^d
- For $P \in \mathcal{H}_k$ we define (kernel of Riesz Transform)

$$K_P(x) = \partial_k \frac{P(x)}{|x|^{k+d}} \quad \text{with} \quad \partial_k = \frac{\Gamma\left(\frac{k+d}{2}\right)}{\pi^{\frac{d}{2}} \Gamma\left(\frac{k}{2}\right)}$$

- when $k=1$, $P(x) = x_j$, then K_{x_j} - is the kernel of classical first order Riesz - transform $R_{x_j} = R_j$

Definitions of Riesz transforms -RT

$$R_p^t f(x) = \int_{|y|>t} K_p(y) f(x-y) dy \quad (\text{Truncated RT})$$

$$R_p f(x) = \lim_{t \rightarrow 0} R_p^t f(x) \quad (\text{RT})$$

$$R_p^* f(x) = \sup_{t>0} |R_p^t f(x)| \quad (\text{Maximal RT})$$

A classical result (E.M. Stein 80')

For each $1 < p < \infty$ there exists $C(p)$ independent of dimension d and such that

$$\left\| \left(\sum_{i=1}^d |R_{x_j} f|^2 \right)^{\frac{1}{2}} \right\|_p \leq C(p) \|f\|_p$$

Why care? Consequences:

$$\| |\nabla f| \|_p \leq C_p \| (-\Delta)^{\frac{1}{2}} f \|_p$$

$$\left\| \left(\sum_{j,k=1}^d |\partial_j \partial_k f|^2 \right)^{\frac{1}{2}} \right\|_p \leq D_p \| \Delta f \|_p$$

Our goals

— Prove a similar result to Stein with R_{x_j} replaced

by $R_{x_j}^*$, e.g.

$$\left\| \left(\sum |R_{x_j}^* f|^2 \right)^{\frac{1}{2}} \right\|_p \leq C_p \|f\|_p$$

— Control $\|R_P^* f\|_p$ in terms of $\|R_P f\|_p$

in a dimension-free fashion

— Dependence on k -degree of P is allowed

— Get explicit dependence in terms of $p \in (1, \infty)$

Previous results

— case $k=1$, $1 < p < \infty$

$$\|R_{x_j}^* f\|_p \leq C(p, d) \|R_{x_j} f\|_p, \quad \text{Mateu, Verdern 2006}$$

Here $C(p, d) \approx_p \frac{1}{d}$

— case $k=1$, $p=2$

$$\|R_{x_j}^* f\|_2 \leq 2 \cdot 10^8 \|R_{x_j} f\|_2 \quad \text{Kucharski, Wróbel 2021}$$

— general odd k , $P \in \mathcal{H}_k$, $1 < p < \infty$

$$\|R_p^* f\|_p \leq C(p, k, d) \|R_p f\|_p$$

Mateu, Orobitg,
Pérez, Verdera 2010

$C(p, k, d) \approx p^{1/k} \gamma^d \rightarrow$ general singular integrals with
odd kernels

— general even k , $P \in \mathcal{H}_k$, $1 < p < \infty$

$$\|R_p^* f\|_p \leq C(p, k, d) \|R_p f\|_p$$

Mateu, Orobitg, Verdera
2011

$C(p, k, d) \approx p^{1/k} \gamma^d \rightarrow$ general singular integrals
with even kernels

Our results: (Kucharski, W, Zienkiewicz 2022, 2023)

- For single Riesz transform

THEOREM 1

Let $k \in \mathbb{N}$. For $1 < p < \infty$ we have

$$\|R_p^* f\|_p \leq C(k) \left(\max(p, (p-1)^{-1})\right)^{2+\frac{k}{2}} \|R_p f\|_p,$$

where $C(k)$ depends only on k but not on the dimension d

— For vectors of Riesz transforms

THEOREM 2

Let $k \in \mathbb{N}$, and let \mathcal{G}_k be a subset of \mathcal{M}_k .

Then for $1 < p < \infty$ we have

$$\left\| \left(\sum_{P \in \mathcal{G}_k} |R_P^* f|^2 \right)^{\frac{1}{2}} \right\|_p \leq C(k) \left(\max_{P \in \mathcal{G}_k} (p, (p-1)^{-1}) \right)^{3 + \frac{k}{2}} \times \\ \times \left\| \left(\sum_{P \in \mathcal{G}_k} |R_P f|^2 \right)^{\frac{1}{2}} \right\|_p .$$

Here $C(k)$ depends only on k , and not on d

Possible approaches - case $k=1$

— Cotlar's inequality

$$|R_{x_j}^* f(x)| \leq C(d) (M(R_{x_j} f)(x) + Mf(x))$$

Problems: $C(d)$ depends on d . Extra term

$Mf(x)$ - Hardy-Littlewood maximal function

— Method of rotations as in Duong-Rubio de Francia

$$R_{x_j}^t f(x) \approx d^{\frac{1}{2}} \int_{S^{d-1}} H_\omega^t f(x) \omega_j d\omega$$

Is dimension-free & works for $\|R_{x_j}^*\|_p \leq C_p \|f\|_p$

we lose $R_{x_j} f(x)$

Our approach - case $k=1$ - TWO STEPS

1. FACTORIZATION

$$R_{x_j}^t = M^t R_{x_j}$$

where $M^t = M_1^t$ - operators with radial kernel

2. METHOD OF ROTATIONS

$$R_{x_j}^t f(x) \approx \sqrt{d} \int_{S^{d-1}} \omega_j H_{\omega}^t f(x) d\omega$$

where H_{ω}^t - directional truncated Hilbert transform

Sketch of TWO STEPS

1. FACTORIZATION

— (Mateu, Orobitz, Pérez, Verdara 2011) (implicitly) proved that

$$R_{x_j}^t = M^t R_{x_j}, \quad j=1, \dots, d$$

$(M^t f)^\wedge(\xi) = m(t\xi) \hat{f}(\xi)$, m -radial and bounded

M^t - bounded on L^p , $1 < p < \infty$.

— Control of $R_{x_j}^*$ in terms, of R_{x_j} is reduced to bounding

$$M^* f(x) = \sup_{t>0} |M^t f(x)|$$

— For single Riesz transform we are left with

$$\|M^* f\|_p \leq C \left(\max (p, (p-1)^{-1}) \right)^{\frac{5}{2}} \|f\|_p$$

Where $1 < p < \infty$, and C is independent of the dimension d

— M^t has a useful explicit formula

$$M^t R_{x_j} R_{x_j} = R_{x_j}^t R_{x_j}$$

$$M^t \sum_{j=1}^d (R_{x_j})^2 = \sum_{j=1}^d R_{x_j}^t R_{x_j}$$

$$M^t = - \sum_{j=1}^d R_{x_j}^t R_{x_j}$$

because $\sum_{j=1}^d (R_{x_j})^2 = -\text{Id}$

— caveat: this does not work for $k > 1$.

— A curiosity (Liu, Melentijević, Zhu 2023)

$$M^t f = \varphi_t * f$$

Where the kernel φ_t is

— non-negative and radial

$$— \|\varphi_t\|_{L^1} = 1$$

— $\varphi_t(x)$ increases for $|x| < t$ and decreases for
for $|x| > t$

— $\varphi_t(x) = \infty$ for $|x| = t$

2. METHOD OF ROTATIONS. (Sketch)

- We focus on the estimate for single Ricci transform
- The goal is to bound $\|M^* f\|_p$. We do not focus on dependence on p . Only on independence of d
- By method of rotations and factorization

$$M^t f = - \sum_{j=1}^d R_j^t R_j f \approx - \sqrt{d} \int_{S^{d-1}} H_\omega^t \left(\sum_j \omega_j R_j f \right) d\omega$$

— Minkowski's inequality gives

$$\|M^* f\|_p \lesssim \sqrt{d} \int_{S^{d-1}} \|H_\omega^* (\sum_j \omega_j R_j f)\|_p d\omega$$

— Because H_ω^* is one-dimensional after Hölder and reverse Hölder on S^{d-1} we get

$$\|M^* f\|_p \leq C_p \left\| \left(\sum_j |R_j f|^2 \right)^{\frac{1}{2}} \right\|_p$$

— By the classical result of Stein

$$\left\| \left(\sum_j |R_j f|^2 \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^d)} \leq C_p \|f\|_{L^p(\mathbb{R}^d)}$$

We are done

Remark: A similar, but more complicated argument works for the vectorial estimate. Extra ingredient: Khintchine inequality.

CASE $k > 1$, odd — THREE STEPS

1. FACTORIZATION, $\rho \in H_k$

$$Q_p^t = M_k^t Q_p$$

where $M^t = M_k^t$ — operators with radial kernel

2. AVERAGING

3. Only then METHOD OF ROTATIONS

Extra ingredient for $k > 1$ — AVERAGING

— Why is it needed?

Say $k=3$. Then

$$\underline{I} = \sum_{i=1}^d \sum_{j=1}^d \sum_{l=1}^d R_{x_i}^2 R_{x_j}^2 R_{x_l}^2$$

But a number of $R_{x_i} R_{x_j} R_{x_l}$ are not higher order even — transforms...

For instance $R_{x_1} R_{x_1} R_{x_1}$ would correspond to $P(x) = x_1^3$
which is not in \mathcal{H}_3 (not harmonic)

— Moreover, the FACTORIZATION formula

$$R_{x_i x_j x_l}^t = M_3^t (R_{x_i} R_{x_j} R_{x_l})$$

works only when i, j, l - pairwise distinct

— How to remedy the problem?

— average over the $SO(d)$ group.

Denote $I = \{(i, j, \ell) \in \{1, \dots, d\}^3 : \text{pairwise distinct}\}$. Then

$$I_d = C(d, u) \int_{SO(d)} \mathcal{E}_u \left(\sum_{n \in I} \left(\mathbb{P}_{x_{n_1}, x_{n_2}, x_{n_3}} \right)^2 \right) (x) d\nu(u)$$

where $C(d, u) \approx 1$, uniformly in d , $d\nu$ - Haar

measure on $SO(d)$, \mathcal{E}_u - conjugation by $u \in SO(d)$

— This procedure in effect reduces our task to controlling the maximal function for

$$R^t f = \sum_{n \in \mathbb{I}} R_{x_{n_1} x_{n_2} x_{n_3}}^t R_{x_{n_1} x_{n_2} x_{n_3}}$$

because M^t has a radial symbol and

$$M^t \left(\sum_{n \in \mathbb{I}} \left(R_{x_{n_1} x_{n_2} x_{n_3}} \right)^2 \right) = \sum_{n \in \mathbb{I}} M^t \left(R_{x_{n_1} x_{n_2} x_{n_3}} \right)^2 = R^t$$

— In the case $k=1$ we had $R^t = M^t$

Case $k > 1$ - even. FIVE STEPS

1. FACTORIZATION
2. AVERAGING
3. EXTENSION TO \mathbb{C}^d
4. COMPLEX METHOD OF ROTATIONS
(IWANIEC AND MARTIN)
5. RESTRICTION TO \mathbb{R}^d

Why are extra steps needed?

- (Real) method of rotations requires that the kernel K_p^t or K_p is odd
- When k is even the kernel is even
- A curiosity: only easy case $k=2$, then

$M_2^t =$ Mandy-Littlewood evenizing operator

Extra ingredients for k -even

STEP 3 Extension to \mathbb{C}^d

— replace K_p^t and K_p - functions on \mathbb{R}^d
with their appropriate complexifications

\tilde{K}_p^t and \tilde{K}_p - functions on $\mathbb{C}^d \cong \mathbb{R}^{2d}$

— for instance

$$\tilde{K}_p(x+iy) = \tilde{Z}_k \frac{P(x+iy)}{|x+iy|^{2d+k}} \quad \tilde{Z}_k = \frac{\Gamma(\frac{k}{2} + d)}{\pi^d \Gamma(\frac{k}{2})}$$

STEP 4. Complex method of rotations

$$\widehat{R}_p^t \tilde{f} = \widehat{K}_p *_{\mathbb{R}^{2d}} \tilde{f} \approx d^{\frac{k}{2}} \int_{S^{2d-1}} P(\omega + i\sigma) H_{\sum_{1 \leq k}^t} \tilde{f} \, d\zeta,$$

— $\zeta = \omega + i\sigma \in \mathbb{C}^d$

— $d\zeta$ — normalized surface measure on S^{2d-1}

— $H_{\Sigma, k}^t$ — k -th power of the complex Hilbert transform truncated at $t > 0$

$$H_{\Sigma, k}^t f(z) = \int_{\mathbb{C}} 1_{|\lambda| \geq t} \left(\frac{\lambda}{|\lambda|} \right)^k \frac{1}{|\lambda|^2} f(z - \lambda \zeta) d\lambda$$

— $H_{\Sigma, k}^t$ is a two-dimensional truncated singular integral operator

STEP 5

Restriction to \mathbb{R}^d

- there is a standard procedure to go back to \mathbb{R}^d
→ integration of $\tilde{K}_p^t(x+iy)$ over $y \in \mathbb{R}^d$
- Problem: we do not get back to $K_p^t(x)$...
Instead we obtain a different kernel $\mathcal{K}_p^t(x)$
- Fortunately their difference can be estimated

General singular integrals?

What about general singular integrals with even or odd kernels Ω ? This is *more subtle ...* If

$$T_{\Omega} f(x) = \int_{\mathbb{R}^d} \frac{\Omega\left(\frac{y}{|y|}\right)}{|y|^{d+1}} f(x-y) dy, \quad T_{\Omega}^* f(x) = \sup_{t>0} \left| \int_{|x-y|>t} \frac{\Omega\left(\frac{y}{|y|}\right)}{|y|^{d+1}} f(x-y) dy \right|$$

Then $\|T_{\Omega}^* f\|_2 \leq C \|T_{\Omega} f\|_2$ may only hold if Ω satisfies an algebraic condition related to its expansion in spherical harmonics (Mateu, Orobitg, Pérez, Verdera 2010, 2011)

Thank you for your attention!